arXiv:1303.2861v1 [math.PR] 12 Mar 2013

Fractional discrete processes: compound and mixed Poisson representations

Luisa Beghin* Claudio Macci†

Abstract

We consider two fractional versions of a family of nonnegative integer valued processes. We prove that their probability mass functions solve fractional Kolmogorov forward equations, and we show the overdispersion of these processes. As particular examples in this family, we can define fractional versions of some processes in the literature as the Polya-Aeppli, the Poisson Inverse Gaussian and the Negative Binomial. We also define and study some more general fractional versions with two fractional parameters.

AMS Subject Classification: 26A33; 33E12; 60G22.
Keywords: Cox process; doubly stochastic Poisson process; Negative Binomial process; Polya Aeppli process; Poisson Inverse Gaussian process.

1 Introduction

In this paper we consider a large family of nonnegative integer valued processes $\{M(t) : t \geq 0\}$ defined by

$$M(t) := \sum_{k=1}^{N_\lambda(t)} X_k, \tag{1}$$

where $\{X_n : n \geq 1\}$ is a sequence of i.i.d. positive and integer valued random variables, independent of a (non-fractional) Poisson process $\{N_\lambda(t) : t \geq 0\}$ with intensity λ. Throughout this paper we also deal with a mixed Poisson representation of the process in (1); more precisely we mean $\{N_1(S(t)) : t \geq 0\}$, where $\{S(t) : t \geq 0\}$ is a subordinator, independent of a (non-fractional) Poisson process $\{N_1(t) : t \geq 0\}$ with intensity 1, such that $N_1(S(t))$ is distributed as $M(t)$ for each fixed $t > 0$.

Our aim is to present and analyze two fractional versions of the process $\{M(t) : t \geq 0\}$ (see **(FV1)** and **(FV2)** at the beginning of Section 3). This allows to define fractional generalizations of some processes in the literature, which include the Polya-Aeppli, the Poisson Inverse Gaussian and the Negative Binomial. These processes are commonly used when the empirical count data exhibit overdispersion, i.e. when the sample variance is larger than the sample mean; moreover it is known that mixed Poisson processes provide simple counting models with overdispersion. We shall see that the fractional versions have the same feature.

The two fractional versions are obtained by considering independent random time-changes of $\{M(t) : t \geq 0\}$ in terms of a stable subordinator $\mathcal{A}^\alpha$ or its inverse $\mathcal{L}^\alpha$ (for $\alpha \in (0, 1)$). This approach is inspired by the recent increasing interest on random time-changed and subordinated processes; see, among the others, [31], [22], [7], [18] and [16]. These processes are widely studied and applied,

---

*Address: Dipartimento di Scienze Statistiche, Sapienza Università di Roma, Piazzale Aldo Moro 5, I-00185 Roma, Italy. e-mail: `luisa.beghin@uniroma1.it`

†Dipartimento di Matematica, Università di Roma Tor Vergata, Via della Ricerca Scientifica, I-00133 Roma, Italia. E-mail: `macci@mat.uniroma2.it`

mainly because their finite dimensional distributions display power law decay and thus heavy tails, either in the case of renewals and birth processes (see [20] and [25]) or in that of diffusions (see [1], [29] and [24]). We recall that fractional versions of the compound Poisson process (as the ones in this paper) have been studied in [32] and [5], in the case of continuous distributed summands.

In this paper we provide a natural fractional extension of some results in the literature. In particular Proposition 3.1 below shows that the probability mass functions of the two fractional versions of the process in (1) solve suitable Kolmogorov equations, where the classical derivatives are replaced by the fractional derivative in the Caputo sense (for the first version), and by the right sided fractional Riemann-Liouville derivatives on $\mathbb{R}_+$ (for the second version).

We conclude with the outline of the paper. We start with some preliminaries in Section 2. In Section 3 we illustrate some properties of the fractional versions of the process in (1). Some examples are presented in Section 4. Finally Section 5 is devoted to more general fractional versions of some processes in Section 4 (the Polya Aeppli and the Poisson Inverse Gaussian) with two fractional parameters.

2 Preliminaries

We start with some preliminaries on fractional calculus, i.e. we give the definitions of two fractional derivatives for real functions defined on $[0,\infty)$.

1. If $\nu \in (0,1)$, the Caputo derivative of order ν (see e.g. (2.4.17) in [11] with $a=0$) is defined by
$$ {}_C D^{\nu}_{0+,t} f(t) := \frac{1}{\Gamma(1-\nu)} \int_0^t (t-s)^{-\nu} \frac{d}{ds} f(s) ds \text{ (for all } t \geq 0). $$
Remark. If we set $\nu = 1$, ${}_C D^{\nu}_{0+,t}$ coincides with the classical derivative $\frac{d}{dt}$; see e.g. Theorem 2.1(b) in [11].

2. If $\nu \in (1,\infty)$, the right sided fractional Riemann-Liouville derivative on $\mathbb{R}_+$ of order ν (see e.g. (2.2.4) in [11]) is defined by
$$ {}_{RL} D^{\nu}_{-,t} f(t) := \frac{1}{\Gamma(m-\nu)} \left(-\frac{d}{dt}\right)^m \int_t^{\infty} \frac{f(s)}{(s-t)^{\nu-m+1}} ds \text{ (for all } t \geq 0), \text{ where } m := \lfloor \nu \rfloor + 1. $$
Remark. If we set $\nu = 1$, ${}_{RL} D^{\nu}_{-,t}$ coincides with $-\frac{d}{dt}$ (i.e. the opposite of the classical derivative); see e.g. (2.2.5) in [11].

Hereafter, for simplicity, we always write
$$ \frac{d^{\nu}_C}{dt^{\nu}} := {}_C D^{\nu}_{0+,t} \text{ (for } \nu \in (0,1)) \text{ and } \frac{d^{\nu}_{RL}}{dt^{\nu}} := {}_{RL} D^{\nu}_{-,t} \text{ (for } \nu \in (1,\infty)). $$

Throughout the paper we often use the symbol $Z(\cdot)$ to mean a process $\{Z(t) : t \geq 0\}$. In view of what follows we recall some preliminaries on the stable subordinator $\mathcal{A}^{\alpha}(\cdot)$ of order $\alpha \in (0,1)$, and its inverse $\mathcal{L}^{\alpha}(\cdot)$. More precisely let $\mathcal{A}^{\alpha}(\cdot)$ be the Lévy process (starting at the origin) such that, for each fixed $t > 0$, we have
$$ \mathbb{E}[e^{\theta \mathcal{A}^{\alpha}(t)}] = \begin{cases} \exp(-(-\theta)^{\alpha} t) & \text{if } \theta \leq 0 \\ \infty & \text{if } \theta > 0; \end{cases} \tag{2} $$
thus, by referring to [30], the random variable $\mathcal{A}^{\alpha}(t)$ has stable distribution of index α and parameters $\mu = 0$, $\theta = 1$ and $\sigma = (t\cos(\frac{\pi\alpha}{2}))^{1/\alpha}$. Furthermore $\mathcal{L}^{\alpha}(\cdot)$ is defined by
$$ \mathcal{L}^{\alpha}(t) := \inf\{z : \mathcal{A}^{\alpha}(z) > t\}; $$

then we have $P(\mathcal{L}^\alpha(t) \leq z) = P(\mathcal{A}^\alpha(z) \geq t)$ for all $z, t > 0$ and

$$\mathbb{E}[e^{-\theta\mathcal{L}^\alpha(t)}] = E_{\alpha,1}(-\theta t^\alpha) \text{ for all } \theta \geq 0. \tag{3}$$

Now we recall the definition of the two following fractional Poisson processes (see e.g. [21] for $\nu \in (0,1)$ and [26] for the case $\nu \in (1,\infty)$). In both cases we consider a random time-change of a non-fractional Poisson process $N_\lambda(\cdot)$.

1. For $\nu \in (0,1)$, $N_\lambda^\nu(\cdot) := N_\lambda(\mathcal{L}^\nu(\cdot))$ where $N_\lambda(\cdot)$ and $\mathcal{L}^\nu(\cdot)$ are independent.
2. For $\nu \in (1,\infty)$, $\hat{N}_\lambda^\nu(\cdot) := N_\lambda(\mathcal{A}^{1/\nu}(\cdot))$ where $N_\lambda(\cdot)$ and $\mathcal{A}^{1/\nu}(\cdot)$ are independent.

We can give some further details and we need to introduce some notation. For all integer $r \geq 0$ and for all $\gamma \in \mathbb{R}$, the rising factorial, also called Pochhammer symbol, is defined by

$$(\gamma)^{(r)} := \begin{cases} \gamma(\gamma+1)\cdots(\gamma+r-1) & \text{if } r \geq 1 \\ 1 & \text{if } r = 0, \end{cases}$$

the falling factorial is defined by

$$(\gamma)_r := \begin{cases} \gamma(\gamma-1)\cdots(\gamma-(r-1)) & \text{if } r \geq 1 \\ 1 & \text{if } r = 0, \end{cases}$$

and we also consider the notation

$$\binom{\gamma}{r} := \begin{cases} \frac{(\gamma)_r}{r!} = \frac{\gamma(\gamma-1)\cdots(\gamma-r+1)}{r!} & \text{if } r \geq 1 \\ 1 & \text{if } r = 0. \end{cases}$$

1. If $\nu \in (0,1)$, it is known (see [21]) that $N_\lambda^\nu(t)$ is distributed as $\sum_{n\geq 1} 1_{\{T_1+\cdots+T_n \leq t\}}$, where $\{T_n : n \geq 1\}$ are i.i.d. random variables with Mittag-Leffler distribution, i.e. with continuous density

   $$f(t) = \lambda t^{\nu-1} E_{\nu,\nu}(-\lambda t^\nu) 1_{(0,\infty)}(t)$$

   (see e.g. [3] and [19]), where $E_{\alpha,\beta}(x) := \sum_{r\geq 0} \frac{x^r}{\Gamma(\alpha r+\beta)}$ is the Mittag-Leffler function (see e.g. [28], page 17). Then, if we consider the generalized Mittag-Leffler function $E_{\alpha,\beta}^\gamma(x) := \sum_{r\geq 0} \frac{(\gamma)^{(r)} x^r}{r!\Gamma(\alpha r+\beta)}$, we have (see formula (2.5) in [4])

   $$P(N_\lambda^\nu(t) = k) = (\lambda t^\nu)^k E_{\nu,\nu k+1}^{k+1}(-\lambda t^\nu) \text{ for all integer } k \geq 0. \tag{4}$$

2. If $\nu \in (1,\infty)$, we have

   $$P(\hat{N}_\lambda^\nu(t) = k) = \frac{(-1)^k}{k!} \sum_{r=0}^\infty \frac{(-\lambda^{1/\nu} t)^r}{r!} (r/\nu)_k \text{ for all integer } k \geq 0. \tag{5}$$

   We recall that this process is presented in [26] by referring to the fractional difference operator $(1-B)^\alpha$, for $\alpha \in (0,1]$, and it is called space-fractional Poisson process. More precisely the probability mass function in (5), with $\nu = 1/\alpha$, satisfies the following equations:

   $$\begin{cases} \frac{d}{dt} P(\hat{N}_\lambda^{1/\alpha}(t) = k) = -\lambda^\alpha (1-B)^\alpha P(\hat{N}_\lambda^{1/\alpha}(t) = k) \text{ for all integer } k \geq 0 \\ P(\hat{N}_\lambda^{1/\alpha}(0) = 0) = 1,\ P(\hat{N}_\lambda^{1/\alpha}(0) = k) = 0 \text{ for all integer } k \geq 1, \end{cases} \tag{6}$$

   where B is the so called *backward shift operator* defined by $Bf(k) = f(k-1)$ and $B^{r-1}Bf(k) = f(k-r)$; thus in particular we have

   $$(1-B)^\alpha f(k) = \sum_{j=0}^\infty (-1)^j \binom{\alpha}{j} f(k-j).$$

   We remark that formula (5) here coincides with formula (1.2) in [26], though it is written in a slightly different way.

3 The fractional versions: compound and mixed representations

We start with the *compound representation* of the fractional versions of the process $M(\cdot)$ in (1), and we refer to the processes defined in Section 2:

(FV1): $M^{\nu}(t) := M(\mathcal{L}^{\nu}(t)) = \sum_{k=1}^{N_{\lambda}^{\nu}(t)} X_k$, for $\nu \in (0,1)$;

(FV2): $\hat{M}^{\nu}(t) := M(\mathcal{A}^{1/\nu}(t)) = \sum_{k=1}^{\hat{N}_{\lambda}^{\nu}(t)} X_k$, for $\nu \in (1,\infty)$.

In view of what follows we consider the following notation for the probability mass function and the probability generating function of the random variables $\{X_n : n \geq 1\}$:

$$q_k := P(X_1 = k) \text{ (for all integer } k \geq 1); \quad g_q(u) := \sum_{k=1}^{\infty} u^k q_k.$$

We also consider the notation

$$q_k^{*n} := P(X_1 + \cdots + X_n = k), \tag{7}$$

and therefore we have $q_k^{*n} = 0$ for all integer $k < n$.

Now we introduce the two following probability mass functions:

1. for $\nu \in (0,1)$, we have

$$p_k^{\nu}(t) := P(M_{\lambda}^{\nu}(t) = k) = \begin{cases} P(N_{\lambda}^{\nu}(t) = 0) & \text{if } k = 0 \\ \sum_{n=1}^{k} q_k^{*n} P(N_{\lambda}^{\nu}(t) = n) & \text{if } k \geq 1; \end{cases} \tag{8}$$

2. for $\nu \in (1,\infty)$, we have

$$\hat{p}_k^{\nu}(t) := P(\hat{M}_{\lambda}^{\nu}(t) = k) = \begin{cases} P(\hat{N}_{\lambda}^{\nu}(t) = 0) & \text{if } k = 0 \\ \sum_{n=1}^{k} q_k^{*n} P(\hat{N}_{\lambda}^{\nu}(t) = n) & \text{if } k \geq 1. \end{cases} \tag{9}$$

The aim of this section is to prove that these probability mass functions satisfy suitable versions of Kolmogorov equations with fractional derivatives.

Proposition 3.1. *If $\nu \in (0,1)$, the probability mass function in (8) satisfies the equations*

$$\begin{cases} \frac{d_C^{\nu}}{dt^{\nu}} p_0^{\nu}(t) = -\lambda p_0^{\nu}(t) \\ \frac{d_C^{\nu}}{dt^{\nu}} p_k^{\nu}(t) = -\lambda p_k^{\nu}(t) + \lambda \sum_{i=1}^{k} q_i p_{k-i}^{\nu}(t) \text{ for all integer } k \geq 1, \end{cases} \tag{10}$$

with the initial conditions $p_0^{\nu}(0) = 1$ and $p_k^{\nu}(0) = 0$ for all integer $k \geq 1$.
If $\nu \in (1,\infty)$, the probability mass function in (9) satisfies the equations

$$\begin{cases} \frac{d_{RL}^{\nu}}{dt^{\nu}} \hat{p}_0^{\nu}(t) = \lambda \hat{p}_0^{\nu}(t) \\ \frac{d_{RL}^{\nu}}{dt^{\nu}} \hat{p}_k^{\nu}(t) = \lambda \hat{p}_k^{\nu}(t) - \lambda \sum_{i=1}^{k} q_i \hat{p}_{k-i}^{\nu}(t) \text{ for all integer } k \geq 1, \end{cases} \tag{11}$$

with the initial conditions $\hat{p}_0^{\nu}(0) = 1$ and $\hat{p}_k^{\nu}(0) = 0$ for all integer $k \geq 1$.
Remarks. *For $\nu = 1$, both formulas (10) and (11) reduce to the well-known Kolmogorov forward equations (for instance one can specialize eq. (2.3) in Chapter 14 in [10]). For $k = 0$, the equations in (10) and (11) coincide with the well-known fractional relaxation equations (see, for example, [2]). For $k > 0$, they can be seen as a discrete version of the fractional master equation (see eq. (5.14) in [21] for the case where the common distribution of the jumps have continuous density; see also [8]).*

Proof. In both cases ($\nu \in (0,1)$ and $\nu \in (1,\infty)$) the initial conditions trivially hold.
Case $\nu \in (0,1)$. We have to check the equations in (10). It is known (see e.g. Theorem 2.1 in [4]) that

$$\left\{\begin{array}{l} \frac{d_C^\nu}{dt^\nu}P(N_\lambda^\nu(t)=0) = -\lambda P(N_\lambda^\nu(t)=0) \\ \frac{d_C^\nu}{dt^\nu}P(N_\lambda^\nu(t)=k) = -\lambda P(N_\lambda^\nu(t)=k) + \lambda P(N_\lambda^\nu(t)=k-1) \text{ for all integer } k \geq 1, \end{array}\right. \tag{12}$$

with the initial conditions

$$P(N_\lambda^\nu(0)=0)=1 \text{ and } P(N_\lambda^\nu(0)=k)=0 \text{ for all integer } k\geq 1.$$

The initial conditions for $(P(N_\lambda^\nu(0)=k))_{k\geq 0}$ meet the ones for $(p_k^\nu(t))_{k\geq 0}$ in the statement of the proposition. For $k=0$ we have $\frac{d_C^\nu}{dt^\nu}p_0^\nu(t) = -\lambda p_0^\nu(t)$ because $p_0^\nu(t) = P(N_\lambda^\nu(t)=0)$. For $k\geq 1$ we have several steps. Firstly, since

$$p_k^\nu(t) = \sum_{n=1}^{h} q_k^{*n} P(N_\lambda^\nu(t)=n) \text{ for all integer } h \geq k,$$

by (12) we get

$$\begin{aligned} \frac{d_C^\nu}{dt^\nu}p_k^\nu(t) =& \sum_{n=1}^{k} q_k^{*n}[-\lambda P(N_\lambda^\nu(t)=n) + \lambda P(N_\lambda^\nu(t)=n-1)] \\ =& -\lambda \sum_{n=1}^{k} q_k^{*n} P(N_\lambda^\nu(t)=n) + \lambda \sum_{n=1}^{k}\left(\sum_{i=1}^{k} q_{k-i}^{*(n-1)} q_i\right) P(N_\lambda^\nu(t)=n-1) \\ =& -\lambda p_k^\nu(t) + \lambda \sum_{i=1}^{k} q_i \sum_{n=1}^{k} q_{k-i}^{*(n-1)} P(N_\lambda^\nu(t)=n-1); \end{aligned}$$

moreover, since $q_h^{*0} = 1_{\{h=0\}}$ and $q_0^{*h} = 1_{\{h=0\}}$, we get

$$\begin{aligned} \sum_{i=1}^{k} q_i \sum_{n=1}^{k} q_{k-i}^{*(n-1)} P(N_\lambda^\nu(t)=n-1) =& \sum_{i=1}^{k-1} q_i \sum_{n=1}^{k} q_{k-i}^{*(n-1)} P(N_\lambda^\nu(t)=n-1) \\ &+ q_k \sum_{n=1}^{k} q_0^{*(n-1)} P(N_\lambda^\nu(t)=n-1) \\ =& \sum_{i=1}^{k-1} q_i \sum_{n=2}^{k} q_{k-i}^{*(n-1)} P(N_\lambda^\nu(t)=n-1) + q_k P(N_\lambda^\nu(t)=0) \\ =& \sum_{i=1}^{k-1} q_i \sum_{j=1}^{k-1} q_{k-i}^{*j} P(N_\lambda^\nu(t)=j) + q_k p_0^\nu(t) \\ =& \sum_{i=1}^{k-1} q_i p_{k-i}^\nu(t) + q_k p_0^\nu(t) = \sum_{i=1}^{k} q_i p_{k-i}^\nu(t), \end{aligned}$$

and this completes the proof.
Case $\nu \in (1,\infty)$. We have to check the equations in (11) and we follow the same lines of the previous case. The main difference concerns only the initial step, i.e. the analogue of (12) presented above. Thus we only give some details on how to prove that we have the following equations

$$\left\{\begin{array}{l} \frac{d_{RL}^\nu}{dt^\nu}P(\hat{N}_\lambda^\nu(t)=0) = \lambda P(\hat{N}_\lambda^\nu(t)=0) \\ \frac{d_{RL}^\nu}{dt^\nu}P(\hat{N}_\lambda^\nu(t)=k) = \lambda P(\hat{N}_\lambda^\nu(t)=k) - \lambda P(\hat{N}_\lambda^\nu(t)=k-1) \text{ for all integer } k \geq 1, \end{array}\right.$$

with the initial conditions

$$P(\hat{N}_\lambda^\nu(0)=0)=1 \text{ and } P(\hat{N}_\lambda^\nu(0)=k)=0 \text{ for all integer } k\geq 1.$$

Thus, since $\hat{N}_\lambda^\nu(\cdot)=N_\lambda(\mathcal{A}^{1/\nu}(\cdot))$, we have

$$P(\hat{N}_\lambda^\nu(t)=k)=\int_0^\infty \frac{(\lambda x)^k}{k!}e^{-\lambda x}f_\nu(x,t)dx \text{ for all integer } k\geq 0,$$

where $f_\nu(x,t):=f_{\mathcal{A}^{1/\nu}(t)}(x)$ is the density of the random variable $\mathcal{A}^{1/\nu}(t)$. Then we get (the first equality holds by eq. (5.17)-(5.18) in [5] with $\gamma=\frac{1}{\nu}$)

$$\begin{aligned}\frac{d_{RL}^\nu}{dt^\nu}P(\hat{N}_\lambda^\nu(t)=k)=&\int_0^\infty \frac{(\lambda x)^k}{k!}e^{-\lambda x}\frac{\partial}{\partial x}f_\nu(x,t)dx=\left[\frac{(\lambda x)^k}{k!}e^{-\lambda x}f_\nu(x,t)\right]_{x=0}^{x=\infty}\\ &-\int_0^\infty\left(-\lambda P(N_\lambda(x)=k)+\lambda P(N_\lambda(x)=k-1)\right)f_\nu(x,t)dx\\ =&\lambda P(\hat{N}_\lambda^\nu(t)=k)-\lambda P(\hat{N}_\lambda^\nu(t)=k-1),\end{aligned}$$

and the proof is complete by taking into account $P(\hat{N}_\lambda^\nu(t)=-1)=0$ for the case $k=0$. □

Remark. If $\nu\in(1,\infty)$, we can also obtain the alternative equation

$$\frac{d}{dt^\nu}\hat{p}_k^\nu(t)=-\lambda^{1/\nu}\sum_{j=0}^k(-1)^j\binom{1/\nu}{j}\sum_{l=0}^\infty q_k^{*(l+j)}P(\hat{N}_\lambda^\nu(t)=l) \text{ for all integer } k\geq 1,$$

by taking the classical derivative with respect to t in $\hat{p}_k^\nu(t)=\sum_{n=1}^k q_k^{*n}P(\hat{N}_\lambda^\nu(t)=n)$, and by (6). It is easy to check that, for $\nu=1$, we meet the classical Kolmogorov equation.

Now let $N_1(\cdot)$ be a non-fractional Poisson process with intensity 1, independent of $\mathcal{L}^\nu(\cdot)$ (for $\nu\in(0,1)$) and of $\mathcal{A}^{1/\nu}(\cdot)$ (for $\nu\in(1,\infty)$). Moreover let $S(\cdot)$ be a subordinator, independent of all the other processes such that the following conditions hold.

1. If $\nu\in(0,1)$, $M^\nu(t)$ is distributed as $N_1(S(\mathcal{L}^\nu(t)))$; in what follows we set $S^\nu(\cdot):=S(\mathcal{L}^\nu(\cdot))$, and we have

$$p_k^\nu(t)=\mathbb{E}\left[\frac{(S^\nu(t))^k}{k!}e^{-S^\nu(t)}\right]=\mathbb{E}\left[\frac{(S(\mathcal{L}^\nu(t)))^k}{k!}e^{-S(\mathcal{L}^\nu(t))}\right] \text{ for all integer } k\geq 0. \quad (13)$$

2. If $\nu\in(1,\infty)$, $\hat{M}^\nu(t)$ is distributed as $N_1(S(\mathcal{A}^{1/\nu}(t)))$; in what follows we set $\hat{S}^\nu(\cdot):=S(\mathcal{A}^{1/\nu}(\cdot))$, and we have

$$\hat{p}_k^\nu(t)=\mathbb{E}\left[\frac{(\hat{S}^\nu(t))^k}{k!}e^{-\hat{S}^\nu(t)}\right]=\mathbb{E}\left[\frac{(S(\mathcal{A}^{1/\nu}(t)))^k}{k!}e^{-S(\mathcal{A}^{1/\nu}(t))}\right] \text{ for all integer } k\geq 0. \quad (14)$$

In such a case we talk about of *mixed representation* of the fractional versions of the process $M(\cdot)$ in (1). Furthermore, in view of what follows, it is useful to introduce the function κ_S defined by $\kappa_S(\theta):=\log\mathbb{E}[e^{\theta S(1)}]$ for all $\theta\in\mathbb{R}$ (note that $\kappa_S(\cdot)$ is a nondecreasing function and $\kappa_S(0)=0$). This function has a strict relationship with the probability generating function g_q presented above; actually, by considering standard computations on independent random time-changes for Lévy processes (see e.g. [6]) one can easily check that

$$\mathbb{E}\left[e^{\theta S(t)}\right]=e^{\lambda t(g_q(1+\theta)-1)} \quad (15)$$

for all θ such that $1+\theta$ belongs to the domain of convergence of g_q.

We conclude with the following further formulas: if $\nu \in (0,1)$, by (3) we have

$$\mathbb{E}\left[e^{-\theta S^\nu(t)}\right] = E_{\nu,1}(\kappa_S(-\theta)t^\nu) \text{ if } \theta \geq 0; \tag{16}$$

if $\nu \in (1,\infty)$, by (2) we have

$$\mathbb{E}\left[e^{\theta \hat{S}^\nu(t)}\right] = \begin{cases} \exp(-(-\kappa_S(\theta))^{1/\nu}t) & \text{if } \theta \leq 0 \\ \infty & \text{if } \theta > 0. \end{cases} \tag{17}$$

On the concept of overdispersion. It is well-known that a real valued random variable Y is said to be overdispersed if $\mathrm{Var}[Y] - \mathbb{E}[Y] > 0$; similarly, for a process $Y(\cdot)$, we have overdispersion if all the random variables $\{Y(t) : t > 0\}$ are overdispersed. Typically the compound Poisson process $M(\cdot)$ in (1) exhibits overdispersion when we exclude the trivial case where the random jumps $\{X_n : n \geq 1\}$ are all equal to 1; actually, in such a case, we have $M(\cdot) = N_\lambda(\cdot)$, i.e. $M(\cdot)$ is a non-fractional Poisson process, and we have $\mathrm{Var}[M(t)] - \mathbb{E}[M(t)] = 0$ for all $t > 0$.

Here we want to study the same feature for the process $M^\nu(\cdot)$ in **(FV1)**; we do not deal with $\hat{M}^\nu(\cdot)$ in **(FV2)** because each random variable $\hat{M}^\nu(t)$ has infinite mean (actually the moment generating function of $\mathcal{A}^{1/\nu}(t)$ is not finite in a neighborhood of the origin; see formula (2) presented above). It is known (see e.g. formulas (2.7) and (2.8) in [3]) that

$$\begin{cases} \mathbb{E}[N_\lambda^\nu(t)] = \frac{\lambda t^\nu}{\Gamma(\nu+1)} \\ \mathrm{Var}[N_\lambda^\nu(t)] = \frac{\lambda t^\nu}{\Gamma(\nu+1)} + (\lambda t^\nu)^2 Z(\nu), \text{ where } Z(\nu) := \frac{1}{\nu}\left(\frac{1}{\Gamma(2\nu)} - \frac{1}{\nu\Gamma^2(\nu)}\right). \end{cases}$$

Thus, since we have $Z(\nu) > 0$ for all $\nu \in (0,1)$ and $Z(1) = 0$ (see Figure 1), the fractional Poisson process $N_\lambda^\nu(\cdot)$ (for $\nu \in (0,1)$) exhibits overdispersion.



Figure 1: Plot of $Z(\nu) := \frac{1}{\nu}\left(\frac{1}{\Gamma(2\nu)} - \frac{1}{\nu\Gamma^2(\nu)}\right)$ versus ν.

Furthermore, it is known (see e.g. Appendix B.4 in [12]) that

$$\begin{cases} \mathbb{E}[M^\nu(t)] = \mathbb{E}[N_\lambda^\nu(t)]\mathbb{E}[X_1] \\ \mathrm{Var}[M^\nu(t)] = \mathbb{E}[N_\lambda^\nu(t)]\mathrm{Var}[X_1] + \mathrm{Var}[N_\lambda^\nu(t)]\mathbb{E}^2[X_1]; \end{cases}$$

then, with some computations, we get

$$\begin{aligned} \mathrm{Var}[M^\nu(t)] - \mathbb{E}[M^\nu(t)] =& \mathbb{E}[N_\lambda^\nu(t)](\mathrm{Var}[X_1] - \mathbb{E}[X_1]) + \mathrm{Var}[N_\lambda^\nu(t)]\mathbb{E}^2[X_1] \\ =& \frac{\lambda t^\nu}{\Gamma(\nu+1)}(\mathbb{E}[X_1^2] - \mathbb{E}[X_1]) + (\lambda t^\nu)^2 Z(\nu)\mathbb{E}^2[X_1] \end{aligned}$$

and $\mathbb{E}[X_1^2] - \mathbb{E}[X_1] \geq 0$ because $P(X_1 \geq 1) = 1$. In conclusion the compound process $M^\nu(\cdot)$ in **(FV1)** exhibits overdispersion.

4 Examples for the process $M(\cdot)$

In this section we present some examples for the process $M(\cdot)$ in (1) which allow to define the corresponding fractional versions $M^\nu(\cdot)$ in **(FV1)** and $\hat{M}^\nu(\cdot)$ in **(FV2)**. More precisely we consider suitable choices of $(q_k)_{k\geq 1}$ and λ; moreover, in all cases, the probability mass function $(q_k)_{k\geq 1}$ is a *zero truncated negative binomial distribution* as in Example B.3.1.5 in Appendix B.3.1 in [12], where β in that reference stands for $\frac{1-\alpha}{\alpha}$, for some $\alpha \in (0,1)$, and $r > -1$:

	$r \neq 0$	$r = 0$ (the limit as $r \to 0$)
probability mass function	$q_k = \frac{\binom{r+k-1}{k}(1-\alpha)^k}{\alpha^{-r}-1}$	$q_k = -\frac{(1-\alpha)^k}{k \log \alpha}$
probability generating function	$g_q(u) = \frac{\alpha^r}{1-\alpha^r}\frac{1-(1-u(1-\alpha))^r}{(1-u(1-\alpha))^r}$	$g_q(u) = \frac{\log(1-u(1-\alpha))}{\log \alpha}$

(where $|u| < 1/(1-\alpha)$ for the probability generating functions). In our examples we have $r = 1$ (geometric distribution) for the Polya Aeppli process, $r = -\frac{1}{2}$ (extended truncated negative binomial distribution) for the Poisson Inverse Gaussian process, and $r = 0$ (logarithmic distribution) for the Negative Binomial process.

Example 1 (Polya Aeppli process). *We choose*

$$\boxed{\alpha = 1-p \text{ and } r = 1 \text{ for some } p \in (0,1) \text{ and } \lambda > 0.}$$

The probability mass functions for the two fractional Polya Aeppli processes can be obtained from (8)*,* (9)*,* (4) *and* (5) *with* $q_k^{*n} = \binom{k-1}{n-1}(1-p)^n p^{k-n}$ *(see e.g. Example 3.19 in [12], where* β *in that reference stands for* $\frac{p}{1-p}$*; another interesting reference on this process is [23]).*
Alternative formulas for the mixed representation can be obtained from (13) *and* (14) *by considering* $S(\cdot)$ *as a compound Poisson process with exponentially distributed summands; more precisely, for* $\mu = \lambda/p$ *and* $\beta = (1-p)/p$*, we mean* $S(\cdot) = \sum_{k=1}^{N_\mu(\cdot)} Y_k^{(\beta)}$*, where* $\{Y_k^{(\beta)} : n \geq 1\}$ *are i.i.d. random variables with density* $f(x) = \beta e^{-\beta x} 1_{(0,\infty)}(x)$*, independent of the (non-fractional) Poisson process* $N_\mu(\cdot)$ *with intensity* μ *(see e.g. Subsection 11.1.2 in [9], which concerns a slightly more general situation with the generalized Polya Aeppli distribution and the Tweedie distribution; in particular we recover our case with the Tweedie distribution with parameters* $(-1, \frac{\lambda(1-p)}{p^2}, \frac{1-p}{p})$*).*

Then we can prove the following result for Example 1 as a consequence of Proposition 3.1.

Proposition 4.1. *Let us consider the probability mass functions in* (8) *and* (9)*, where* $M(\cdot)$ *as in Example 1. If* $\nu \in (0,1)$*, we have*

$$\begin{cases} \frac{d_C^\nu}{dt^\nu} p_0^\nu(t) = -\lambda p_0^\nu(t) \\ \frac{d_C^\nu}{dt^\nu} p_k^\nu(t) - (1-\alpha)\frac{d_C^\nu}{dt^\nu} p_{k-1}^\nu(t) = -\lambda p_k^\nu(t) + \lambda p_{k-1}^\nu(t) \text{ for all integer } k \geq 1 \end{cases} \tag{18}$$

with the initial conditions $p_0^\nu(0) = 1$ *and* $p_k^\nu(0) = 0$ *for all integer* $k \geq 1$*.*
If $\nu \in (1,\infty)$*, we have*

$$\begin{cases} \frac{d_{RL}^\nu}{dt^\nu} \hat{p}_0^\nu(t) = \lambda \hat{p}_0^\nu(t) \\ \frac{d_{RL}^\nu}{dt^\nu} \hat{p}_k^\nu(t) - (1-\alpha)\frac{d_{RL}^\nu}{dt^\nu} \hat{p}_{k-1}^\nu(t) = \lambda \hat{p}_k^\nu(t) - \lambda \hat{p}_{k-1}^\nu(t) \text{ for all integer } k \geq 1 \end{cases} \tag{19}$$

with the initial conditions $\hat{p}_0^\nu(0) = 1$ *and* $\hat{p}_k^\nu(0) = 0$ *for all integer* $k \geq 1$*.*
Remark. *The equations in* (18) *and* (19) *coincide for* $\nu = 1$*. They are equations for the probability mass functions of the non-fractional Polya Aeppli process, and we do not know any references where they appear.*

Proof. Firstly, for $k = 0$, we have the same equations in Proposition 3.1; thus we restrict the attention on the case $k \geq 1$. If $\nu \in (0,1)$, by (10) and by taking into account that $q_h = (1-\alpha)^{h-1}\alpha$ (for all integer $h \geq 1$), we have

$$\begin{aligned}\frac{d_C^\nu}{dt^\nu}p_k^\nu(t) - (1-\alpha)\frac{d_C^\nu}{dt^\nu}p_{k-1}^\nu(t) =& -\lambda p_k^\nu(t) + \lambda\sum_{i=1}^{k}(1-\alpha)^{i-1}\alpha p_{k-i}^\nu(t)\\ &-(1-\alpha)\left(-\lambda p_{k-1}^\nu(t) + \lambda\sum_{i=1}^{k-1}(1-\alpha)^{i-1}\alpha p_{k-1-i}^\nu(t)\right)\\ =& -\lambda p_k^\nu(t) + \lambda\alpha p_{k-1}^\nu(t) + \lambda\sum_{i=2}^{k}(1-\alpha)^{i-1}\alpha p_{k-i}^\nu(t)\\ &+\lambda(1-\alpha)p_{k-1}^\nu(t) - \lambda\sum_{i=1}^{k-1}(1-\alpha)^{i}\alpha p_{k-1-i}^\nu(t)\\ =& -\lambda p_k^\nu(t) + \lambda p_{k-1}^\nu(t).\end{aligned}$$

If $\nu \in (1,\infty)$, the proof follows the same lines; we have to consider (11) instead of (10), and there are suitable changes of signs. □

Example 2 (Poisson Inverse Gaussian process)**.** *We have*

$$\boxed{\alpha = 1 - \tfrac{2\beta}{1+2\beta},\ r = -\tfrac{1}{2} \text{ and } \lambda = \lambda_{\beta,\mu} := \tfrac{\mu}{\beta}((1+2\beta)^{1/2}-1) \text{ for some } \beta,\mu > 0.}$$

We are not aware of any reference with a formula for the convolution densities q_k^{*n} *in* (7)*. So* (8) *and* (9) *do not give completely explicit formulas for the probability mass functions of the two fractional Poisson Inverse Gaussian processes.*

The formulas (13) *and* (14) *for the mixed representation can be obtained by considering* $S(\cdot)$ *as an Inverse Gaussian process; more precisely* $S(t) = Y_{\mu,\beta}(t)$ *should have density*

$$f(x) = \frac{\mu t}{(2\pi\beta x^3)^{1/2}}\exp\left(-\frac{(x-\mu t)^2}{2\beta x}\right)1_{(0,\infty)}(x). \tag{20}$$

Moreover (see e.g. eq. (3.39) in [12]) we have

$$\mathbb{E}\left[e^{\theta Y_{\mu,\beta}(t)}\right] = \begin{cases}\exp\left(-\frac{\mu}{\beta}t\left((1-2\beta\theta)^{1/2}-1\right)\right) & \text{if } \theta < \frac{1}{2\beta}\\ \infty & \text{if } \theta \geq \frac{1}{2\beta}\end{cases} \tag{21}$$

We remark that the processes $S^\nu(\cdot)$ (if $\nu \in (0,1)$) and $\hat{S}^\nu(\cdot)$ (if $\nu \in (1,\infty)$) for Example 2 can be seen as a fractional Inverse Gaussian process. A different fractional version of this process has been defined in [15] by subordinating a fractional Brownian motion to an Inverse Gaussian process (in analogy with the so-called fractional Laplace motion; see [13]). Some properties of the process defined in [15] are illustrated in [17].

Example 3 (Negative Binomial process)**.** *We choose*

$$\boxed{\alpha = p,\ r = 0 \text{ and } \lambda = -\log p \text{ for some } p \in (0,1).}$$

The probability mass functions for the two fractional Negative Binomial processes can be obtained from (8)*,* (9)*,* (4) *and* (5) *with* $q_k^{*n} = \frac{n!}{(-\log p)^n}\frac{(1-p)^k|s(k,n)|}{k!}$*, where* $\{|s(k,n)| : k \in \{0,1,\dots,n\}\}$ *are the* unsigned Stirling numbers of the first kind *(see e.g. Theorem 6 in [27]).*

Alternative formulas for the mixed representation can be obtained from (13) *and* (14) *by considering* $S(\cdot)$ *as a Gamma process; more precisely* $S(t)$ *should have density*

$$f(x) = \frac{(\frac{p}{1-p})^t}{\Gamma(t)}x^{t-1}e^{-\frac{p}{1-p}x}1_{(0,\infty)}(x).$$

5 Processes with two fractional parameters

In this section we generalize the fractional versions of the processes in Examples 1 and 2 by considering a further fractional parameter $\eta \in (0,1)$. Throughout this section we use the notation $N_1^{(\eta)}(\cdot)$ for the process $N_1^{1/\eta}(\cdot)$ in (5); then, for each example, we consider independent random time-changes of $N_1^{(\eta)}(\cdot)$ in terms of the processes $S^\nu(\cdot)$ in (13) and $\hat{S}^\nu(\cdot)$ in (14), i.e. the fractional versions of the subordinator $S(\cdot)$ which appears in the mixed representation of $M^\nu(\cdot)$ presented in Section 3 above. Thus we have

$$p_k^{\eta,\nu}(t) := P(M^{\eta,\nu}(t) = k), \text{ where } M^{\eta,\nu}(\cdot) = N_1^{(\eta)}(S^\nu(\cdot)), \text{ if } \nu \in (0,1); \tag{22}$$

$$\hat{p}_k^{\eta,\nu}(t) := P(\hat{M}^{\eta,\nu}(t) = k), \text{ where } \hat{M}^{\eta,\nu}(\cdot) = N_1^{(\eta)}(\hat{S}^\nu(\cdot)), \text{ if } \nu \in (1,\infty). \tag{23}$$

We remark that we recover the cases presented above by setting $\eta = 1$. We shall see that the governing equations for these probability mass functions have not only a fractional time derivative but also the fractional difference operator $(1-B)^\eta$ as (6).

5.1 Generalized fractional Polya Aeppli process

We start with the result.

Proposition 5.1. *Let $S(\cdot) = \sum_{k=1}^{N_\mu(\cdot)} Y_k^{(\beta)}$ be as in Example 1. If $\nu \in (0,1)$, the probability mass function in (22) satisfies the equations*

$$\begin{cases} \frac{d_C^\nu}{dt^\nu} p_0^{\eta,\nu}(t) = -\frac{1}{\beta}\left(\mu + \frac{d_C^\nu}{dt^\nu}\right) p_0^{\eta,\nu}(t) \\ \frac{d_C^\nu}{dt^\nu} p_k^{\eta,\nu}(t) = -\frac{1}{\beta}\left(\mu + \frac{d_C^\nu}{dt^\nu}\right)(1-B)^\eta p_k^{\eta,\nu}(t) \text{ for all integer } k \geq 1, \end{cases} \tag{24}$$

with the initial conditions $p_0^{\eta,\nu}(0) = 1$ and $p_k^{\eta,\nu}(0) = 0$ for all integer $k \geq 1$.
If $\nu \in (1,\infty)$, the probability mass function in (23) satisfies the equations

$$\begin{cases} \frac{d_{RL}^\nu}{dt^\nu} \hat{p}_0^{\eta,\nu}(t) = \frac{1}{\beta}\left(\mu - \frac{d_{RL}^\nu}{dt^\nu}\right) \hat{p}_0^{\eta,\nu}(t) \\ \frac{d_{RL}^\nu}{dt^\nu} \hat{p}_k^{\eta,\nu}(t) = \frac{1}{\beta}\left(\mu - \frac{d_{RL}^\nu}{dt^\nu}\right)(1-B)^\eta \hat{p}_k^{\eta,\nu}(t) \text{ for all integer } k \geq 1, \end{cases} \tag{25}$$

with the initial conditions $\hat{p}_0^{\eta,\nu}(0) = 1$ and $\hat{p}_k^{\eta,\nu}(0) = 0$ for all integer $k \geq 1$.
Remark. *The equations in (24) and (25) coincide for $\nu = 1$.*

Proof. In both cases ($\nu \in (0,1)$ and $\nu \in (1,\infty)$) the initial conditions trivially hold.
Case $\nu \in (0,1)$. We have

$$P(S^\nu(t) \in G) = P(N_\mu(\mathcal{L}^\nu(t)) = 0)1_G(0) + \int_G f_\nu(x,t)dx \text{ (for all Borel subsets } G \text{ of } [0,\infty)),$$

where f_ν is the density of the absolutely continuous part of the random variable $S^\nu(t)$; thus, since $P(N_\mu(\mathcal{L}^\nu(t)) = 0) = E_{\nu,1}(-\mu t^\nu)$, we obtain

$$p_k^{\eta,\nu}(t) = E_{\nu,1}(-\mu t^\nu)1_{\{k=0\}} + \int_0^\infty P(N_1^{(\eta)}(\mathcal{L}^\nu(z)) = k) f_\nu(z,t)dz.$$

Thus we get

$$\begin{aligned} \frac{d_C^\nu}{dt^\nu} p_k^{\eta,\nu}(t) &= \left(-\mu E_{\nu,1}(-\mu t^\nu) + \frac{1}{\beta}\left(\mu + \frac{d_C^\nu}{dt^\nu}\right) f_\nu(0,t)\right) 1_{\{k=0\}} \\ &\quad + \frac{1}{\beta}\left(\mu + \frac{d_C^\nu}{dt^\nu}\right) \int_0^\infty \frac{d}{dz}\{P(N_1^{(\eta)}(\mathcal{L}^\nu(z)) = k)\} f_\nu(z,t)dz \end{aligned}$$

by (2.4.58) in [11], (2.5) in [5] and an integration by parts (where we take into account that $P(N_1^{(\eta)}(\mathcal{L}^\nu(0))=k)=1_{\{k=0\}}$). The last equation reduces to

$$\frac{d_C^\nu}{dt^\nu}p_k^{\eta,\nu}(t)=\frac{1}{\beta}\left(\mu+\frac{d_C^\nu}{dt^\nu}\right)\int_0^\infty\frac{d}{dz}\{P(N_1^{(\eta)}(\mathcal{L}^\nu(z))=k)\}f_\nu(z,t)dz$$

because

$$-\mu E_{\nu,1}(-\mu t^\nu)+\frac{1}{\beta}\left(\mu+\frac{d_C^\nu}{dt^\nu}\right)f_\nu(0,t)=0. \tag{26}$$

We remark that eq. (26) can be checked by inspection after noting that:

$$f_\nu(0,t)=\mu\beta t^\nu E_{\nu,\nu+1}^2(-\mu t^\nu)=\mu\beta t^\nu\sum_{j=0}^\infty\frac{(j+1)(-\mu t^\nu)^j}{\Gamma(\nu j+\nu+1)}=-\beta\sum_{j=0}^\infty\frac{j(-\mu t^\nu)^j}{\Gamma(\nu j+1)}$$

by eq. (2.4) in [5] and some computations;

$$\frac{d_C^\nu}{dt^\nu}f_\nu(0,t)=\mu\beta\sum_{j=0}^\infty\frac{(j+1)(-\mu)^j\frac{d_C^\nu}{dt^\nu}t^{\nu j+\nu}}{\Gamma(\nu j+\nu+1)}=\mu\beta\sum_{j=0}^\infty\frac{(j+1)(-\mu t^\nu)^j}{\Gamma(\nu j+1)}$$

by taking into account that we have $\frac{d_C^\nu}{dt^\nu}t^{\nu j+\nu}=\frac{\Gamma(\nu j+\nu+1)}{\Gamma(\nu j+1)}t^{\nu j}$ by (2.1.17) in [11].
Finally we can conclude the proof by using (6) with $\lambda=1$, and with some computations (where we distinguish the cases $k=0$ and $k\geq 1$).
Case $\nu\in(1,\infty)$. We have

$$P(\hat{S}^\nu(t)\in G)=P(N_\mu(\mathcal{A}^{1/\nu}(t))=0)1_G(0)+\int_G\hat{f}_\nu(x,t)dx \text{ (for all Borel subsets } G \text{ of } [0,\infty)),$$

where $\hat{f}_\nu$ is the density of the absolutely continuous part of the random variable $\hat{S}^\nu(t)$; thus, since $P(N_\mu(\mathcal{A}^{1/\nu}(t))=0)=e^{-\mu^{1/\nu}t}$, we obtain

$$\hat{p}_k^{\eta,\nu}(t)=e^{-\mu^{1/\nu}t}1_{\{k=0\}}+\int_0^\infty P(N_1^{(\eta)}(\mathcal{A}^{1/\nu}(z))=k)\hat{f}_\nu(z,t)dz.$$

We follow the same lines of the proof for the previous case and we give only some details. Firstly we get

$$\begin{aligned}\frac{d_{RL}^\nu}{dt^\nu}p_k^{\eta,\nu}(t)=&\left(\mu e^{-\mu^{1/\nu}t}-\frac{1}{\beta}\left(\mu-\frac{d_{RL}^\nu}{dt^\nu}\right)f_\nu(0,t)\right)1_{\{k=0\}}\\&-\frac{1}{\beta}\left(\mu-\frac{d_{RL}^\nu}{dt^\nu}\right)\int_0^\infty\frac{d}{dz}\{P(N_1^{(\eta)}(\mathcal{A}^{1/\nu}(z))=k)\}\hat{f}_\nu(z,t)dz.\end{aligned}$$

by (2.2.15) in [11], (5.8) in [5] and an integration by parts (where we take into account that $P(N_1^{(\eta)}(\mathcal{A}^{1/\nu}(0))=k)=1_{\{k=0\}}$). The last equation reduces to

$$\frac{d_{RL}^\nu}{dt^\nu}p_k^{\eta,\nu}(t)=-\frac{1}{\beta}\left(\mu-\frac{d_{RL}^\nu}{dt^\nu}\right)\int_0^\infty\frac{d}{dz}\{P(N_1^{(\eta)}(\mathcal{A}^{1/\nu}(z))=k)\}\hat{f}_\nu(z,t)dz$$

because

$$\mu e^{-\mu^{1/\nu}t}-\frac{1}{\beta}\left(\mu-\frac{d_{RL}^\nu}{dt^\nu}\right)f_\nu(0,t)=0. \tag{27}$$

We remark that eq. (27) can be checked by inspection after noting that: $f_\nu(0,t) = \frac{\beta\mu^{1/\nu}t}{\nu}e^{-\mu^{1/\nu}t}$ by eq. (5.7) in [5];

$$\begin{aligned}\frac{d^\nu_{RL}}{dt^\nu}f_\nu(0,t) =&\frac{\beta\mu^{1/\nu}}{\nu}\frac{d^\nu_{RL}}{dt^\nu}\left\{te^{-\mu^{1/\nu}t}\right\} = \frac{\beta\mu^{1/\nu}}{\nu}\frac{d^\nu_{RL}}{dt^\nu}\left(-\frac{d}{d(\mu^{1/\nu})}e^{-\mu^{1/\nu}t}\right)\\ =&-\frac{\beta\mu^{1/\nu}}{\nu}\frac{d}{d(\mu^{1/\nu})}\frac{d^\nu_{RL}}{dt^\nu}e^{-\mu^{1/\nu}t} = -\frac{\beta\mu^{1/\nu}}{\nu}\frac{d}{d(\mu^{1/\nu})}\left\{(\mu^{1/\nu})^\nu e^{-\mu^{1/\nu}t}\right\}\\ =&-\frac{\beta\mu^{1/\nu}}{\nu}\left(\nu\mu\cdot\mu^{-1/\nu}e^{-\mu^{1/\nu}t}-t\mu e^{-\mu^{1/\nu}t}\right)\end{aligned}$$

by (2.2.15) in [11] (for the equality $\frac{d^\nu_{RL}}{dt^\nu}e^{-\mu^{1/\nu}t} = (\mu^{1/\nu})^\nu e^{-\mu^{1/\nu}t}$) and some computations.
Finally we conclude by using (6) with $\lambda = 1$ and with some computations (where we distinguish the cases $k = 0$ and $k \geq 1$). □

We already know that, for $\eta = 1$, $M^{\eta,\nu}(\cdot)$ and $\hat{M}^{\eta,\nu}(\cdot)$ coincide with the fractional versions in **(FV1)** and **(FV2)** when we deal with the Polya Aeppli process $M(\cdot)$ as in Example 1. Then one can check that the results in Proposition 4.1 can be recovered by considering Proposition 5.1 with $\eta = 1$. Actually $(1-B)^1 p_k^{1,\nu}(t) = p_k^{1,\nu}(t) - p_{k-1}^{1,\nu}(t)$ and, if we rearrange the terms in a suitable way, the equations (24) and (25) with $\eta = 1$ meet (18) and (19), respectively, by setting $\mu = \frac{\lambda}{p} = \frac{\lambda}{1-\alpha}$ and $\beta = \frac{1-p}{p} = \frac{\alpha}{1-\alpha}$ as in Example 1.

5.2 Generalized fractional Poisson Inverse Gaussian process

From now on we consider the usual symbol $\delta(x)$ for the Dirac delta. Furthermore we consider the process $S(\cdot) = Y_{\mu,\beta}(\cdot)$ as in Example 2, and we introduce the following notation.

$$f_\nu(x,t) = f_{S^\nu(t)}(x) \text{ is the density of the random variable } S^\nu(t), \text{ for } \nu \in (0,1); \tag{28}$$

$$\hat{f}_\nu(x,t) = f_{S^\nu(t)}(x) \text{ is the density of the random variable } \hat{S}^\nu(t), \text{ for } \nu \in (1,\infty). \tag{29}$$

The main result in this subsection is Proposition 5.3 which provides governing equations for the probability mass functions $(p_k^{\eta,\nu}(t))_{k\geq 0}$ and $(\hat{p}_k^{\eta,\nu}(t))_{k\geq 0}$ when $S(\cdot)$ is as in Example 2. In its proof we refer to Proposition 5.2, i.e. a preliminary result for f_ν and $\hat{f}_\nu$; thus, in some sense, Proposition 5.2 concerns the fractional versions of the Poisson Inverse Gaussian process in Example 2, and not its generalized version considered here.

Proposition 5.2. *Let f_ν and $\hat{f}_\nu$ be the functions in (28) and (29), respectively. If $\nu \in (0,1)$, we have*

$$\frac{\partial^\nu_C}{\partial t^\nu}\frac{\partial^\nu_C}{\partial t^\nu}f_\nu(x,t) - 2\frac{\mu}{\beta}\frac{\partial^\nu_C}{\partial t^\nu}f_\nu(x,t) = 2\frac{\mu^2}{\beta}\frac{\partial}{\partial x}f_\nu(x,t) \tag{30}$$

with the initial conditions $f_\nu(x,0) = \delta(x)$ and $f_\nu(0,t) = 0$.
If $\nu \in (1,\infty)$, we have

$$\frac{\partial^\nu_{RL}}{\partial t^\nu}\frac{\partial^\nu_{RL}}{\partial t^\nu}\hat{f}_\nu(x,t) + 2\frac{\mu}{\beta}\frac{\partial^\nu_{RL}}{\partial t^\nu}\hat{f}_\nu(x,t) = 2\frac{\mu^2}{\beta}\frac{\partial}{\partial x}\hat{f}_\nu(x,t) \tag{31}$$

with the initial conditions $\hat{f}_\nu(x,0) = \delta(x)$ and $\hat{f}_\nu(0,t) = 0$.
Remark. *The equations in (30) and (31) coincide for $\nu = 1$, and coincide with the equation Theorem 3.1 in [15] (with the same initial conditions).*

Proof. We introduce the following notation: $\tilde{g}(\theta)$, for $\theta \geq 0$, is the Laplace transform of $g(x)$; $g^*(\sigma)$, for $\sigma \geq 0$, is the Laplace transform of $g(t)$ (we need the second one only for $\nu \in (0,1)$). Furthermore we recall that, by (21), we have

$$\kappa_S(-\theta) = \log \mathbb{E}[e^{-\theta S(1)}] = -\frac{\mu}{\beta}\left((1+2\beta\theta)^{1/2} - 1\right).$$

Case $\nu \in (0,1)$. We have $\tilde{f}_\nu(\theta,t) = E_{\nu,1}(\kappa_S(-\theta)t^\nu)$ by (16), and

$$\frac{\partial_C^\nu}{\partial t^\nu}\tilde{f}_\nu(\theta,t) = \kappa_S(-\theta)\tilde{f}_\nu(\theta,t) \tag{32}$$

by eq. (2.4.58) in [11]. Moreover we have

$$\left(\frac{\partial_C^\nu}{\partial t^\nu} g\right)^*(\sigma) = \sigma^\nu g^*(\sigma) - \sigma^{\nu-1}g(0) \tag{33}$$

by eq. (2.4.63) in [11]. We take the Laplace transforms with respect to t in (32) and, by taking into account (33) with $g(\cdot) = \tilde{f}_\nu(\theta,\cdot)$ for the left hand side, we get

$$\sigma^\nu \tilde{f}_\nu^*(\theta,\sigma) - \sigma^{\nu-1}\underbrace{\tilde{f}_\nu(\theta,0)}_{=E_{\nu,1}(0)=1} = \kappa_S(-\theta)\tilde{f}_\nu^*(\theta,\sigma);$$

then we have

$$0 = (\sigma^\nu - \kappa_S(-\theta))\tilde{f}_\nu^*(\theta,\sigma) - \sigma^{\nu-1} = \left(\sigma^\nu - \frac{\mu}{\beta} + \frac{\mu}{\beta}(1+2\beta\theta)^{1/2}\right)\tilde{f}_\nu^*(\theta,\sigma) - \sigma^{\nu-1}$$

and, after some computations, we obtain

$$\left(\sigma^\nu\left(\sigma^\nu - 2\frac{\mu}{\beta}\right) - 2\frac{\mu^2}{\beta}\theta\right)\tilde{f}_\nu^*(\theta,\sigma) - \sigma^{\nu-1}\left(\sigma^\nu - 2\frac{\mu}{\beta} + \kappa_S(-\theta)\right) = 0. \tag{34}$$

We complete the proof for the case $\nu \in (0,1)$ by checking that (30) yields (34). This will be done in what follows. We take the Laplace transforms with respect to x in (30) and, after an integration by parts where we take into account that $f_\nu(0,t) = 0$, we have

$$\frac{\partial_C^\nu}{\partial t^\nu}\frac{\partial_C^\nu}{\partial t^\nu}\tilde{f}_\nu(\theta,t) - 2\frac{\mu}{\beta}\frac{\partial_C^\nu}{\partial t^\nu}\tilde{f}_\nu(\theta,t) = 2\frac{\mu^2}{\beta}\theta\tilde{f}_\nu(\theta,t);$$

then we take the Laplace transforms with respect to t and, by taking into account (33) with $g(\cdot) = \frac{\partial_C^\nu}{\partial t^\nu}\tilde{f}_\nu(\theta,\cdot)$ and with $g(\cdot) = \tilde{f}_\nu(\theta,\cdot)$ together with some computations, we get

$$\left(\sigma^\nu - 2\frac{\mu}{\beta}\right)\left(\sigma^\nu\tilde{f}_\nu^*(\theta,\sigma) - \sigma^{\nu-1}\underbrace{\tilde{f}_\nu(\theta,0)}_{=1}\right) - \sigma^{\nu-1}\left.\frac{\partial_C^\nu}{\partial t^\nu}\tilde{f}_\nu(\theta,t)\right|_{t=0} = 2\frac{\mu^2}{\beta}\theta\tilde{f}_\nu^*(\theta,\sigma);$$

finally we meet (34) because we have $\left.\frac{\partial_C^\nu}{\partial t^\nu}\tilde{f}_\nu(\theta,t)\right|_{t=0} = \kappa_S(-\theta)$ (by (32) with $t=0$) and $\tilde{f}_\nu(\theta,0) = 1$.

Case $\nu \in (1,\infty)$. We have $\tilde{\hat{f}}_\nu(\theta,t) = \exp(-(-\kappa_S(-\theta))^{1/\nu}t)$ by (17), and $\frac{\partial_{RL}^\nu}{\partial t^\nu}\tilde{\hat{f}}_\nu(\theta,t) = -\kappa_S(-\theta)\tilde{\hat{f}}_\nu(\theta,t)$ and $\frac{\partial_{RL}^\nu}{\partial t^\nu}\frac{\partial_{RL}^\nu}{\partial t^\nu}\tilde{\hat{f}}_\nu(\theta,t) = (\kappa_S(-\theta))^2\tilde{\hat{f}}_\nu(\theta,t)$ by eq. (2.2.15) in [11]. Then one can obtain the equality

$$\left(\frac{\partial_{RL}^\nu}{\partial t^\nu}\frac{\partial_{RL}^\nu}{\partial t^\nu}\tilde{\hat{f}}_\nu\right)(\theta,t) + 2\frac{\mu}{\beta}\left(\frac{\partial_{RL}^\nu}{\partial t^\nu}\tilde{\hat{f}}_\nu\right)(\theta,t) = 2\frac{\mu^2}{\beta}\theta\tilde{\hat{f}}_\nu(\theta,t) \tag{35}$$

with some computations. We complete the proof showing that (31) yields (35). We take the Laplace transforms with respect to x in (31), and we have

$$\left(\frac{\partial^{\nu}_{RL}}{\partial t^{\nu}}\frac{\partial^{\nu}_{RL}}{\partial t^{\nu}}\tilde{\hat{f}}_{\nu}\right)(\theta,t)+2\frac{\mu}{\beta}\left(\frac{\partial^{\nu}_{RL}}{\partial t^{\nu}}\tilde{\hat{f}}_{\nu}\right)(\theta,t)=\int_0^{\infty}e^{-\theta x}2\frac{\mu^2}{\beta}\frac{\partial}{\partial x}\hat{f}_{\nu}(x,t)dx;$$

then we meet (35) by considering an integration by parts for the right hand side, where we take into account the equality $\hat{f}_{\nu}(0,t)=0$. □

Now we are ready to prove Proposition 5.3.

Proposition 5.3. *Let* $(p_k^{\eta,\nu}(t))_{k\geq 0}$ *and* $(\hat{p}_k^{\eta,\nu}(t))_{k\geq 0}$ *be the probability mass functions in* (22) *and* (23)*, respectively, with* $S(\cdot)=Y_{\mu,\beta}(\cdot)$ *as in Example 2. If* $\nu\in(0,1)$*, we have*

$$\frac{d^{\nu}_{C}}{dt^{\nu}}\frac{d^{\nu}_{C}}{dt^{\nu}}p_k^{\eta,\nu}(t)-2\frac{\mu}{\beta}\frac{d^{\nu}_{C}}{dt^{\nu}}p_k^{\eta,\nu}(t)=2\frac{\mu^2}{\beta}(1-B)^{\eta}p_k^{\eta,\nu}(t)\text{ for all integer }k\geq 0 \tag{36}$$

with the initial conditions $p_0^{\eta,\nu}(0)=1$ *and* $p_k^{\eta,\nu}(0)=0$ *for all integer* $k\geq 1$.
If $\nu\in(1,\infty)$*, we have*

$$\frac{d^{\nu}_{RL}}{dt^{\nu}}\frac{d^{\nu}_{RL}}{dt^{\nu}}\hat{p}_k^{\eta,\nu}(t)+2\frac{\mu}{\beta}\frac{d^{\nu}_{RL}}{dt^{\nu}}\hat{p}_k^{\eta,\nu}(t)=2\frac{\mu^2}{\beta}(1-B)^{\eta}\hat{p}_k^{\eta,\nu}(t)\text{ for all integer }k\geq 0 \tag{37}$$

with the initial conditions $\hat{p}_0^{\eta,\nu}(0)=1$ *and* $\hat{p}_k^{\eta,\nu}(0)=0$ *for all integer* $k\geq 1$.
Remark. *The equations in* (36) *and* (37) *coincide for* $\nu=1$.

Proof. In both cases ($\nu\in(0,1)$ and $\nu\in(1,\infty)$) the initial conditions trivially hold.
We start with the case $\nu\in(0,1)$ and we consider the function f_{ν} in (28). Firstly we have

$$p_k^{\eta,\nu}(t)=\int_0^{\infty}P(N_1^{(\eta)}(z)=k)f_{\nu}(z,t)dz\text{ (for all integer }k\geq 0).$$

Then we get

$$\begin{aligned}\frac{d^{\nu}_{C}}{dt^{\nu}}\frac{d^{\nu}_{C}}{dt^{\nu}}p_k^{\eta,\nu}(t)-2\frac{\mu}{\beta}\frac{d^{\nu}_{C}}{dt^{\nu}}p_k^{\eta,\nu}(t)&=\int_0^{\infty}P(N_1^{(\eta)}(z)=k)\left(\frac{\partial^{\nu}_{C}}{\partial t^{\nu}}\frac{\partial^{\nu}_{C}}{\partial t^{\nu}}f_{\nu}(z,t)-2\frac{\mu}{\beta}\frac{\partial^{\nu}_{C}}{\partial t^{\nu}}f_{\nu}(z,t)\right)dz\\&=\int_0^{\infty}P(N_1^{(\eta)}(z)=k)2\frac{\mu^2}{\beta}\frac{\partial}{\partial z}f_{\nu}(z,t)dz\end{aligned}$$

by (30) in Proposition 5.2; then

$$\begin{aligned}\frac{d^{\nu}_{C}}{dt^{\nu}}\frac{d^{\nu}_{C}}{dt^{\nu}}p_k^{\eta,\nu}(t)-2\frac{\mu}{\beta}\frac{d^{\nu}_{C}}{dt^{\nu}}p_k^{\eta,\nu}(t)=&-2\frac{\mu^2}{\beta}\int_0^{\infty}\frac{d}{dz}\{P(N_1^{(\eta)}(z)=k)\}f_{\nu}(z,t)dz\\=&2\frac{\mu^2}{\beta}(1-B)^{\eta}\underbrace{\int_0^{\infty}P(N_1^{(\eta)}(z)=k)f_{\nu}(z,t)dz}_{=p_k^{\eta,\nu}(t)}\end{aligned}$$

by an integration by parts and by taking into account $f_{\nu}(0,t)=0$ (for the first equality) and by (6) with $\lambda=1$ (for the second equality). Thus (36) holds and this completes the proof for the case $\nu\in(0,1)$. The proof for the case $\nu\in(1,\infty)$ follows the same lines; we get (37) by considering a suitable change of sign and, in particular, (31) instead of (30). □

We already know that, for $\eta=1$, $M^{\eta,\nu}(\cdot)$ and $\hat{M}^{\eta,\nu}(\cdot)$ coincide with the fractional versions in **(FV1)** and **(FV2)** when we deal with the Poisson Inverse Gaussian process $M(\cdot)$ as in Example

2. Then one can check that, for $k = 0$, the equations in Proposition 5.3 can be obtained from the equations in Proposition 3.1 adapted to this process. Actually, for $\nu \in (0, 1)$, we have

$$\frac{d_C^\nu}{dt^\nu} p_0^{1,\nu}(t) = -\lambda_{\beta,\mu} p_0^{1,\nu}(t) = -\frac{\mu}{\beta}((1 + 2\beta)^{1/2} - 1) p_0^{1,\nu}(t)$$

by (10) in Proposition 3.1; then we obtain

$$\frac{d_C^\nu}{dt^\nu} \frac{d_C^\nu}{dt^\nu} p_0^{1,\nu}(t) - 2\frac{\mu}{\beta} \frac{d_C^\nu}{dt^\nu} p_0^{1,\nu}(t) = 2\frac{\mu^2}{\beta} p_0^{1,\nu}(t)$$

with some computations, and this meets (36) in Proposition 5.3 because $(1 - B)^1 p_k^{1,\nu}(t) = p_k^{1,\nu}(t) - p_{k-1}^{1,\nu}(t)$ and $p_{-1}^{1,\nu}(t) = 0$ for all $t \geq 0$. For $\nu \in (1, \infty)$ we have similar computations with suitable changes of signs (we have (11) and (37) in place of (10) and (36), respectively).

Acknowledgements. We would like to thank the referee for the careful reading of the paper and for useful suggestions which led to an improved presentation. We also thank Barbara Pacchiarotti for Figure 1.

References

[1] H. Allouba, W. Zheng. Brownian-time processes: the PDE connection and the half-derivative generator. *Ann. Probab.* **29** (2001) 1780–1795.

[2] L. Beghin. Fractional relaxation equations and Brownian crossing probabilities of a random boundary. *Adv. in Appl. Probab.* **44** (2012) 479–505.

[3] L. Beghin, E. Orsingher. Fractional Poisson processes and related planar motions. *Electron. J. Probab.* **14** (2009) 1790–1827.

[4] L. Beghin, E. Orsingher. Poisson-type processes governed by fractional and higher-order recursive differential equations. *Electron. J. Probab.* **15** (2010) 684–709.

[5] L. Beghin, C. Macci. Alternative forms of compound fractional Poisson processes. *Abstr. Appl. Anal.* **2012**, Article ID 747503, 30 pp.

[6] J. Bertoin. *Lévy Processes.* Cambridge University Press, Cambridge, 1996.

[7] C.C. Heyde, N.N. Leonenko. Student processes. *Adv. in Appl. Probab.* **37** (2005) 342–365.

[8] R. Hilfer, L. Anton. Fractional master equations and fractal random walks. *Phys. Rev. E* **51** (1995) R848–R851.

[9] N.L. Johnson, A.W. Kemp, S. Kotz. *Univariate Discrete Distributions (3rd edition).* John Wiley and Sons, Hoboken, NJ, 2005.

[10] S. Karlin, H.M. Taylor. *A Second Course in Stochastic Processes.* Academic Press, London, 1981.

[11] A.A. Kilbas, H.M. Srivastava, J.J. Trujillo. *Theory and Applications of Fractional Differential Equations.* Elsevier, Amsterdam, 2006.

[12] S.A. Klugman, H.H. Panjer, G.E. Willmot. *Loss Models. From Data to Decisions.* John Wiley and Sons, New York, 1998.

[13] T.J. Kozubowski, M.M. Meershaert, K. Podgórski. Fractional Laplace motion. *Adv. in Appl. Probab.* **38** (2006) 451–464.

[14] T.J. Kozubowski, K. Podgórski. Distributional properties of the Negative Binomial Lévy process. *Probab. Math. Statist.* **29** (2009) 43–71.

[15] A. Kumar, M.M. Meerschaert, P. Vellaisamy. Fractional normal inverse Gaussian diffusion. *Statist. Probab. Lett.* **81** (2011) 146–152.

[16] A. Kumar, E. Nane, P. Vellaisamy. Time-changed Poisson processes. *Statist. Probab. Lett.* **81** (2011) 1899–1910.

[17] A. Kumar, P. Vellaisamy. Fractional normal inverse Gaussian process. *Methodol. Comput. Appl. Probab.* **14** (2012) 263–283.

[18] W. Linde, Z. Shi. Evaluating the small deviation probabilities for subordinated Lévy processes. *Stochastic Process. Appl.* **113** (2004) 273–287.

[19] F. Mainardi, R. Gorenflo, E. Scalas. A fractional generalization of the Poisson processes. *Vietnam J. Math.* **32** (2004) 53–64.

[20] F. Mainardi, R. Gorenflo, A. Vivoli. Renewal processes of Mittag-Leffler and Wright type. *Fract. Calc. Appl. Anal.* **8** (2005) 7–38.

[21] M.M. Meerschaert, E. Nane, P. Vellaisamy. The fractional Poisson process and the inverse stable subordinator. *Electron. J. Probab.* **16** (2011) 1600–1620.

[22] M.M. Meerschaert, H.P. Scheffler. Limit theorems for continuous-time random walks with infinite mean waiting times. *J. Appl. Probab.* **41** (2004) 623–638.

[23] L.D. Minkova. The Polya-Aeppli process and ruin problems. *J. Appl. Math. Stoch. Anal.* **2004** (2004) 221-234.

[24] E. Orsingher, L. Beghin. Fractional diffusion equations and processes with randomly varying time. *Ann. Probab.* **37** (2009) 206–249.

[25] E. Orsingher, F. Polito. Fractional pure birth processes. *Bernoulli* **16** (2010) 858–881.

[26] E. Orsingher, F. Polito. The space-fractional Poisson process. *Statist. Probab. Lett.* **82** (2012) 852–858.

[27] G.P. Patil, J.K. Wani. On certain structural properties of the logarithmic series distribution and the first type Stirling distribution. *Sankhyā Ser. A* **27** (1965) 271–280.

[28] I. Podlubny. *Fractional Differential Equations.* Academic Press, San Diego, 1999.

[29] A.I. Saichev, G.M. Zaslawsky. Fractional kinetic equations: solutions and applications. *Chaos* **7** (1997) 753–764.

[30] G. Samorodnitsky, M.S. Taqqu. *Stable Non-Gaussian Random Processes.* Chapman Hall, New York, 1994.

[31] K. Sato. Subordination and self-decomposability. *Statist. Probab. Lett.* **54** (2001) 317–324.

[32] E. Scalas. A class of CTRWs: compound fractional Poisson processes. In: *Fractional Dynamics* (J. Klafter, S.C. Lim and R. Metzler Eds.), pp. 353–374. World Sci. Publ., Hackensack, NJ, 2012.